

# ***Efficacy of angler catch data as a population and conservation monitoring tool for the flagship Mahseer fishes (*Tor* spp.) of Southern India***

ADRIAN C. PINDER<sup>a,b,\*</sup>, RAJEEV RAGHAVAN<sup>a,c,d</sup> and J. ROBERT BRITTON<sup>b</sup>

<sup>a</sup>*Mahseer Trust, c/o The Freshwater Biological Association, East Stoke River Laboratory, Wareham, Dorset, UK*

<sup>b</sup>*Faculty of Science and Technology, Bournemouth University, Fern Barrow, Poole, Dorset, UK*

<sup>c</sup>*Conservation Research Group (CRG), St Albert's College, Kochi, Kerala, India*

<sup>d</sup>*Laboratory of Systematics, Ecology and Conservation, Zoo Outreach Organization (ZOO), Coimbatore, Tamil Nadu, India*

## ABSTRACT

1. Mahseer (*Tor* spp.) are flagship fishes in South Asian rivers. Their populations are threatened through poaching and habitat disturbance, yet they are highly prized game fishes due to their large size, appearance and sporting qualities. The international recreational angling community has frequently been cited as playing a vital role in conserving these fishes while also providing economic benefit to poor rural communities.

2. Owing to a lack of scientific data and the considerable challenges associated with monitoring fish populations in large monsoonal rivers, efforts to determine the long-term trends in their populations has focused on sport-fishing catch records. Here, catch data collected between 1998 and 2012 from Galibore, a former fishing camp on the River Cauvery, Karnataka, India, were analysed to determine the catch per unit effort (CPUE – by number and weight) as an indicator of relative fish abundance, along with the size structure of catches. This fishery operated a mandatory catch-and-release (C&R) policy, and provided the fish community with protection from illegal fishing.

3. Between 1998 and 2012, 23 620 hours fishing effort were applied to catch and release 6161 mahseer, ranging in size from 1 to 104 lbs (0.45–46.8 kg) in weight. Across the period, CPUE in number increased significantly over time with a concomitant decrease in CPUE by weight, revealing strong recruitment in the population and a shift in population size structure. This suggests a strong response to the C&R policy and the reduction in illegal fishing, indicating that conservation strategies focusing on the beneficial and negative aspects of exploitation can be successful in achieving positive outcomes.

4. These outputs from angler catch data provide insights into the mahseer population that were impossible to collect by any alternative method. They provide the most comprehensive analysis of a long-term dataset specific to any of the mahseer species across their entire geographical range and demonstrate the value of organised angling as a conservation monitoring tool to enhance biological data, and inform conservation and fishery management actions. Copyright © 2014 John Wiley & Sons, Ltd.

Received 28 April 2014; Revised 9 September 2014; Accepted 8 November 2014

KEY WORDS: angler logs; C&R; poaching; Western Ghats; stock protection; IUCN Red List; ecosystem services; population monitoring

\*Correspondence to: Adrian Pinder, Mahseer Trust, c/o The Freshwater Biological Association, East Stoke River Laboratory, Wareham, Dorset, BH20 6BB, UK. E-mail: apinder@bournemouth.ac.uk

## INTRODUCTION

Freshwater fishes comprise one of the most threatened taxa on earth (Cooke *et al.*, 2012; Carrizo *et al.*, 2013; Reid *et al.*, 2013), with the extinction of approximately 60 species since 1500 and a further 1679 currently threatened with extinction (Carrizo *et al.*, 2013). Despite that, conservation attention on these fishes is limited, mostly attributable to issues relating to knowledge gaps on key life-history traits, population and habitat requirements, and geographical distributions, all of which are crucial for developing and implementing effective conservation actions (Cooke *et al.*, 2012). Moreover, these knowledge gaps are increasing as taxonomists continue to discover and describe new species of freshwater fishes, many of them from habitats that are already facing high levels of human disturbance.

Collection of inland fisheries data, particularly in biodiversity-rich, tropical countries, can be extremely challenging as many of the sites are located in remote areas and extreme habitats that are often inaccessible for research, and where a lack of political will further limits both financial capacity and human resource (Mahon, 1997; Arce-Ibarra and Charles, 2008). Improving knowledge and understanding of freshwater fish and inland fisheries in these countries and regions therefore needs to consider the use of alternative, cost-effective approaches (Bene *et al.*, 2009; Raghavan *et al.*, 2011; De Graaf *et al.*, in press). Owing to the often threatened status of the fish species concerned, allied with legislation that seeks to protect these species (even if management strategies are yet to be developed because of knowledge gaps), these alternative approaches should also be non-destructive and have a strong ethical basis.

Mahseer (*Tor* spp; Cypriniformes: Cyprinidae) are large-bodied freshwater fishes that are endemic to the monsoonal rivers of Asia. They are popular throughout their range as flagship species of considerable economic, recreational and conservation interest (Singh and Sharma, 1998; Siraj *et al.*, 2007; Nguyen *et al.*, 2008). Of the 18 valid species of *Tor* mahseer (Kottelat, 2013;

Eschmeyer, 2014), six species (*Tor ater*, *Tor khudree*, *Tor kulkarnii*, *Tor malabaricus*, *Tor putitora* and *Tor yunnanensis*) are 'Endangered', one is 'Near Threatened', and six species are 'Data Deficient' on the IUCN Red List (IUCN, 2013; www.iucnredlist.org). The remaining five species have not been assessed for their conservation status. Despite this, data on mahseer populations are severely limited, with even fundamental aspects such as taxonomy, autecology, and population demographics being unknown for many species (Raghavan *et al.*, 2011; Pinder and Raghavan, 2013). For example, there are no population estimates available for the endangered species *T. khudree* and *T. malabaricus* (Raghavan, 2011; Raghavan and Ali, 2011). Nevertheless, they are internationally recognised for their large size, attractive appearance, and sporting qualities by recreational anglers; in India, they are known as the 'King of aquatic systems' (Langer *et al.*, 2001; Dhillon, 2004) and comprise one of the primary groups of fish targeted by recreational fishers (Cooke *et al.*, in press). Indeed, the little information that is available on Indian mahseer populations has largely originated from, or is related to, the recreational angling community (Thomas, 1873; MacDonald, 1948; Trans World Fishing Team, 1984; Dhillon, 2004; Everard and Kataria, 2011; Pinder and Raghavan, 2013).

The recreational angling community offers a social group that positively supports fish conservation (Arlinghaus, 2006) and recreational fishers have engaged in various activities contributing to freshwater fish conservation such as monitoring, research, management, advocacy, and education (Granek *et al.*, 2008; Cooke *et al.*, in press). For example, in India, the recreational fishing sector has played an integral part in the conservation and management of mahseers through such activities as the implementation of compulsory catch-and-release (C&R), stock augmentation, stock protection and, in some cases, the maintenance of catch log-books (Everard and Kataria, 2011; Pinder and Raghavan, 2013; Cooke *et al.*, in press). Nevertheless, despite recreational fishers and fishery managers having previously been identified as a potentially valuable source of data, there have been, to date, no previous

efforts to exploit these catch log-books. Consequently, in this study, catch log-book data from the Galibore Fishing Camp on the River Cauvery were assessed over a 15-year period (1998 to 2012). In this period, the fishery management objectives were the release of all rod-caught mahseers and the elimination of poaching throughout the controlled (~7 km) length of river through enforcement. The study objectives were thus: (i) to determine the temporal trends in catch per unit effort (CPUE – by number and weight) of mahseer captured by recreational fishers; (ii) to assess the extent to which the size structure of the mahseer population has changed over time and how this might be related to the fishery management objectives; and (iii) to assess the implications of the outputs in relation to recreational fishery exploitation and species conservation.

**MATERIALS AND METHODS**

The Cauvery (basin area of 87 900 km<sup>2</sup>) (De Silva *et al.*, 2007) is a major east-flowing river draining the Western Ghats, an exceptional area of freshwater biodiversity and endemism in peninsular India (Molur *et al.*, 2011). The Cauvery and its tributaries comprise one of the two

(the other being the Himalayan Ganges) river systems where C&R angling for the mahseer has been practised since colonial times (Thomas, 1873; Dhu, 1923; MacDonald, 1948). Galibore Fishing Camp is one of four former angling camps situated on the River Cauvery encompassed by the Cauvery Wildlife Sanctuary (an IUCN Category IV Protected Area) in the state of Karnataka, southern India (Figure 1). The Wildlife Association of South India (WASI) came into existence in 1972 with a mandate ‘to conserve and preserve the wildlife of South India’. This Bangalore based non-governmental organization was instrumental in the early development of the C&R fishery which encompassed the 7 km beat at Galibore and extended 22 km between Mutthatti and Mekedatu (Figure 1). Owing to the recognized revenue potential of the fishery, Galibore along with two further camps (sited between Galibore and Shivasamudram Falls) were developed in 1999 into semi-permanent eco-tourism establishments by the state government-owned Jungle Lodges and Resorts (JLR). WASI’s successful model of employing guards to man anti-poaching camps was maintained and supported by both WASI and JLR at Galibore until 2012, when the entire fishery was closed (Pinder and Raghavan, 2013).




Figure 1. Location of the River Cauvery and the study area. Solid line represents the 7 km Galibore fishery. The dashed line represents the 22 km C&R fishery formerly controlled by WASI. Locations are coded: SF: Shivasamudram Falls, MU: Mutthattii, G: Galibore, MT: Mekedatu.

Despite present contention regarding the taxonomic identity of mahseer species within this section of the Cauvery, there are two well-defined morphs known as blue-finned mahseer and golden or hump-back mahseer. As work to resolve the exact identity of these 'species' is under way, this paper refers only to the phenotypic descriptions as 'blue-finned' and 'golden' mahseers to avoid risk of perpetuating erroneous scientific names.

The fishing season for mahseer typically extends from November to March, with fishery performance considered to peak, providing consistent sport quality (number and size of fish caught) between January and March when river flows are at their lowest and angling can be practised effectively.

Between mid-January and mid-March of 1998 to 2012 the mahseer fishery was subject to regulated angling pressure (maximum 10 rods per day), practising a very strict C&R policy. Structured catch data collected during this period included daily records of individual angler identity (name); hours fished (effort); number of fish caught; weight of individual fish (the standard metric used by anglers was imperial lbs) and notes relating to mahseer phenotype. With the exception of 1999 and 2000, a sub-sample of catch returns from 1998 to 2012 were available from the fishery manager and complemented by additional returns retained by anglers over the same period. The resolution of the recovered dataset is summarized in Table 1.

While all larger mahseer (>10 lbs (>4.5 kg)) were typically weighed to the nearest pound using spring-loaded weighing scales, many weights of smaller individual fish were found to be restricted to estimates. Furthermore, where an angler recorded a large number of fish during a single (4h) fishing session, records were typically limited to the weight of the largest fish with the remaining catch enumerated, e.g. 'six fish to 18 lbs'. Following consultation with the camp manager and a selection of the anglers, these data were standardized by recording one fish at 18 lbs with all other individuals recorded as weighing 5 lbs (5 lbs representing the threshold at which most anglers were considered neither to weigh nor to estimate the weight of their fish). Where the

Table 1. Temporal resolution of data recovered to inform CPUE. Individual angler numbers per year and hours fished (effort) between January and March 1998–2012

Year	No. hours fished			Total no. anglers	Total no. hours fished
	Jan	Feb	March		
1998		580		6	580
1999					0
2000					0
2001			820	9	820
2002			1080	10	1080
2003		1920		19	1920
2004		1868	772	25	2640
2005	848		1756	28	2604
2006	264	1344		17	1608
2007	976	1656		27	2632
2008	736	2028	424	33	3188
2009	692	504		11	1196
2010	848	1136		29	1984
2011	984	976	428	35	2388
2012	980			10	980

weight of the largest individual did not exceed 5 lbs (either estimated or weighed), e.g. 'nine fish to 5 lbs', data were standardized by applying a 50% weight reduction to the remaining eight fish for which weights were not recorded. In this example the adjusted record would account for one fish of 5 lbs and eight fish of 2.5 lbs. While the authors acknowledge the inherent limitations of these standardized data, the allocation of arbitrary weights (as guided by the local angling community) has facilitated a valuable measure of the numbers of young fish recruiting to the population over the course of the study period.

The initial step in the data analyses was to determine CPUE by number and weight for each year. These data were then analysed in linear mixed models where the final model used angler identity as the random variable (to account in the model for differences in their respective abilities, differences in fishing style, etc., and in relation to their catches), year as the independent variable and CPUE (either in number or weight) as the dependent variable. Outputs included estimated marginal means (i.e. mean adjusted CPUE by year) and the significance of their differences between years according to pairwise comparisons with Bonferroni adjustment for multiple comparisons. In addition, the mean weights of fish captured per year were tested using ANOVA with Tukey's

post hoc tests. All statistics were completed in SPSS v.21.0.

**RESULTS**

Annual median CPUE increased over the period, although the within-year variability of the data was considerable (Figure 2). The linear mixed models were significant for both catch per unit effort by number ( $F_{12,251}=18.56$ ,  $P <0.01$ ) and weight ( $F_{12,251}=6.13$ ,  $P <0.01$ ), with pairwise comparisons having significantly higher CPUE by number between 2010 and 2012 compared with the highest CPUE by number recorded in the early 2000s (2001;  $P <0.01$ ; Figure 3). There were no significant differences, however, in the mean adjusted catch per unit effort by weight per year ( $P >0.05$ ; Figure 3).




Figure 2. Box plot of year versus catch per unit effort (CPUE) of: (a) number of fish per angler per hour, and (b) weight (lbs) of fish per angler per hour, where the median, 25th and 75th percentile, and 10th and 90th percentile are displayed.




Figure 3. Mean adjusted catch per unit effort by number (a) and weight (b) by year, where the random effects of individual anglers in the data set have been accounted for in the model. \* Difference in catch per unit effort is significantly different from the highest value recorded in the early 2000s ( $P <0.01$ ). Error bars represent standard error.

Over the study period, the mean weight of fish captured by anglers significantly decreased (ANOVA,  $F_{12,251}=7.41$ ,  $P <0.01$ ), with Tukey’s post-hoc tests showing that the differences between the highest mean weight recorded in the study, 1998, and subsequent years were significant between 2007 and 2012 ( $P <0.05$ ; Figure 4). The significant relationship between CPUE by number and mean weight of fish showed that as catch rates increased over time they comprised larger numbers of smaller fish (linear regression:  $R^2=0.83$ ,  $F_{1,11}=22.93$ ,  $P <0.01$ ; Figure 4). Indeed, by categorizing the captured fish into weight categories of 20–39 lbs, 40–59 lbs and >60




Figure 4. (a) Mean weight of fish captured per year; \* Difference in mean weight significantly different from highest values in the early 2000s ( $P < 0.01$ ). (b) Relationship of mean adjusted catch per unit effort per year and the mean weight of fish captured in that year. In all cases, error bars represent standard error.

lbs, it was apparent that the contribution of the largest fish to catches significantly reduced between 2001 and 2012 (linear regression:  $R^2 = 0.82$ ,  $F_{1,10} = 18.81$ ,  $P < 0.01$ ; Figure 5), but not in the smaller weight classes (21–40 lbs: linear regression:  $R^2 = 0.12$ ,  $F_{1,10} = 1.21$ ,  $P = 0.47$ ; 41–60 lbs: linear regression:  $R^2 = 0.57$ ,  $F_{1,10} = 0.57$ ,  $P = 0.47$ ; Figure 5).

## DISCUSSION

The Indian Wildlife (Protection) Act 1972 (WPA) was designed to provide much needed legal protection to flora and fauna. Although this piece




Figure 5. Plot of proportion of weight class of fish to total catch per unit effort by number according to year, where white boxes = 20–39 lbs, grey = 40–59 lbs, and black = >60 lbs (1 lb = 0.45 kg).

of legislation prohibits the hunting of any ‘wild animal’ within areas set aside for protection (Protected Areas (PA)), the Act only specifies amphibians, birds, mammals, and reptiles as constituting the term ‘wild animal’ (Pinder and Raghavan, 2013). Lacking any formal amendment to recognize and include freshwater fish, recently revised government interpretation of the Act has resulted in the closure of the four former recreational mahseer fishing camps sited within the Cauvery Wildlife Sanctuary. The phased closure of these camps between 2010 and 2012 has left fish stocks previously afforded protection from poachers once again vulnerable to the effects of illegal and highly destructive harvest methods, including the use of dynamite (Pinder and Raghavan, 2013). Lacking any scientifically derived survey data, the daily catches recorded by anglers at the Galibore Camp between 1998 and 2012 represent the only available data to examine the temporal performance of the mahseer stock leading up to the implementation of the angling ban, and to explore any potential effects that the C&R fishery may have had on the health of the population.

The outputs of the analyses of the catch data from the Galibore fishery showed some marked changes in catch composition over the study period, with increased numbers of smaller fish appearing in catches allied with increased CPUE by number. This successful use of recreational catch data to obtain insights into the mahseer

population mirror other examples of using recreational angler catch data as a tool to monitor freshwater fish stocks and inform population management strategies (Cowx, 1991; Granek *et al.*, 2008). As a consequence of historic overexploitation, examples in many cases relate to species of high economic value, either as food or sport fishes, which are now facing global or local population threats, e.g. Atlantic salmon *Salmo salar* (Gee and Milner, 1980) and white sturgeon *Acipenser transmontanus* (Inglis and Rosenau, 1994). In the case of 'Endangered' species that are endemic to developing countries (e.g. Eurasian taimen *Hucho taimen* (Jensen *et al.*, 2009); mahseer *Tor* spp. (Pinder and Raghavan, 2013)), resources available to monitor and manage fish populations are typically highly constrained. This limits the development of effective management strategies that are urgently required to foster a balance between exploitation and species conservation (Jensen *et al.*, 2009). Thus, angler catch data can constitute a very cost-effective alternative in collating temporal and spatial information on the fish stock, thereby providing information on long-term population patterns and trends in that component of the stock that is being exploited (Cooke *et al.*, in press).

Although bait selection and angling method can be highly selective with respect to species and sizes of fish captured (Mezzera and Largiadèr, 2001; Ussi-Heikkilä *et al.*, 2008), such bias was considered to be minimal here owing to the very large mouth gape of even the smallest mahseers. Despite some limited effort applied by anglers to catch fish using artificial lures, the primary method of capture was to use large balls (~8 cm diameter) of cereal (ragi, *Eleusine coracana*) derived paste as bait that appeared to capture fish between 1 lb and 104 lbs (0.45–46.8 kg). This was likely to have reduced the potential for variability in the data occurring through using different methods. As any inherent variance in individual angler ability in the dataset was also accounted for in the analyses, the increased appearance of smaller fish in catches suggests that this was due to their greater availability to anglers. These data highlight an apparent threshold between 2007 and 2008, when CPUE by fish number and total

weight demonstrated a marked increase. Given that anecdotal evidence has suggested minimal stock augmentation in the river (S. Chakrabarti, Wildlife Association of South India, pers. comm.), the increased abundance of smaller mahseer has been interpreted as occurring through elevated natural recruitment success. The mechanisms responsible for the observed sudden increase in numbers are not yet understood, but the strong year classes observed since 2008 could potentially be explained by several years of more favourable environmental conditions (e.g. flows) being temporally synchronized with key life-history functions (e.g. spawning and early development).

When considering the abundance of fish recorded within weight categories, fish smaller than 20 lbs (<9 kg) were omitted from the analysis to guarantee the exclusion of all weights derived by the standardized assumptions described in the Methods section. Focusing only on fish weights assigned by individual anglers, it was apparent that the contribution of the largest fish (greater than 60 lbs (>27 kg)) to catches significantly reduced between 2001 and 2012 (Figure 5). While this will have contributed to the overall decrease in mean weight over the same period, it is important to note that these larger specimens were represented by a distinct phenotype and referred to by anglers as 'golden' mahseer or the ambiguous '*Tor mussullah*' (Knight *et al.*, 2013; Pinder and Raghavan, 2013). Establishing the true species identity and conservation status of these larger specimens lies beyond the scope of the present study; however, the notes associated with the current dataset indicate the recent (post-2005) failure in recruitment of this golden phenotype. The resolution of data collected by anglers between 1998 and 2012 therefore goes beyond the provision of just numbers and weights and might also contribute a better understanding of conservation ecology in defining the temporal genetic composition of mahseer within this part of the River Cauvery.

Environmental factors also require consideration of their influence on catch statistics. Potential drivers of catch success include river temperature (McMichael and Kaya, 1991), flow (North, 1980), and turbidity (Drenner

*et al.*, 1997; Lehtonen *et al.*, 2009); all of which can vary in response to natural climatic conditions or in response to river regulation and the manipulation of flows from upstream dams and reservoirs (North and Hickley, 1977; Barillier *et al.*, 1993). Although environmental data are not available to complement the present dataset, environmental factors were likely to have played only a minimal role in influencing angling success over the study period owing to the limited intra-annual timeframes of focus (January–March), when weather and river conditions were typically stable as they are outside of the monsoon season.

In a recent review, Cooke *et al.* (in press) highlighted a global interest by recreational anglers in targeting endangered fish and proposed a dichotomous decision tree of indicators to show whether the practice of C&R angling constitutes a conservation problem or conservation action. The data recorded from the Galibore Camp between 1998 and 2012 clearly demonstrate a natural and significant increase in mahseer population size. However, qualifying the efficacy of the C&R management and stock protection programme in driving the observed increase in fish biomass remains constrained by a lack of comparative empirical data from control sites, which were not afforded protection over the same period. There are many references specifically documenting the long-term efforts of the Cauvery fishing camps and the role of WASI in protecting fish stocks by forcing poaching activities beyond the boundaries of the fishery (Nair, 2010; Pinder, 2013; Pinder and Raghavan, 2013). Despite the largely anecdotal nature of this information, the data presented within the present study, coupled with the fact that recreational fishing interest for these highly prized fish has not since shifted beyond the boundaries of the closed fishery, strengthen the evidence supporting the effective conservation benefits of the former management model practised within the wildlife sanctuary.

In light of the consistent fishery management practice applied across all four former camps and throughout the entire controlled reach, it is considered that the Galibore catch data represent the performance of the mahseer population

throughout the 22 km between Mutthatti and Mokedatu Gorge (Figure 1). Within the broader context of catchment management (Nguyen *et al.*, 2008) and associated ecosystem services (Everard, 2013), the population growth and high biomass of mahseer shown to be present until 2012 may also have been significant at the catchment level. Indeed, in addition to the natural dispersal behaviour typically exhibited by rheophilic cyprinids (Robinson *et al.*, 1998; Reichard *et al.*, 2004), annual monsoon river flows are likely to have been highly effective in delivering larvae and juveniles to the downstream reaches where annual augmentation of the stock would have contributed to maintaining local populations and enhancing the harvest potential for sustenance fishers in downstream rural communities.

In summary, this structured catch dataset collected by recreational anglers visiting Galibore between 1998 and 2012 represents the most comprehensive long-term dataset specific to any of the mahseers across their entire geographical range in Asia. It demonstrates also the value of organized angling as a monitoring tool to enhance biological data and inform conservation and fishery management. These data also provide a unique baseline against which the population response to the recent and radical change in management policy and the closure of the catch and release fishery can be compared with future conservation targets.

#### ACKNOWLEDGEMENTS

The authors wish to thank Sportquest Holidays Ltd, D. Plummer and M. Brown for the provision of catch records. Maps and historical insight to the fishery were provided by S. Chakrabarti and S. Raj. We are also grateful to A. Kanagavel for assisting in the production of Figure 1. This research was supported by the Mahseer Trust.

#### REFERENCES

- Arce-Ibarra AM, Charles AT. 2008. Inland fisheries of the Mayan Zone in Quintana Roo, Mexico: using a combined approach to fishery assessment for data-sparse fisheries. *Fisheries Research* **91**: 151–159.

- Arlinghaus R. 2006. Overcoming human obstacles to conservation of recreational fishery resources, with emphasis on central Europe. *Environmental Conservation* **33**: 46–59.
- Barillier A, Garnier J, Coste M. 1993. Experimental reservoir water release: impact on the water quality on a river 60 km downstream (upper Seine river, France). *Water Research* **27**: 635–643.
- Bene C, Abban EK, Zwietaen P, Dankwa HR, Brummett R, Ofori JK, Obirih-Opareh N, Kolding J. 2009. *Engaging Local Communities in Aquatic Resources Research and Activities: a Technical Manual*. WorldFish Center Technical Manual. WorldFish Center: Penang, Malaysia
- Carrizo SF, Smith KG, Darwall WRT. 2013. Progress towards a global assessment of the status of freshwater fishes (Pisces) for the IUCN Red List: application to conservation programmes in zoos and aquariums. *International Zoo Yearbook* **47**: 46–64.
- Cooke SJ, Paukert C, Hogan Z. 2012. Endangered river fish: factors hindering conservation and restoration. *Endangered Species Research* **17**: 171–191.
- Cooke SJ, Hogan ZS, Butcher PA, Stokesbury MJW, Raghavan R, Gallagher AJ, Hammerschlag N, Danylchuk AJ. in press. Angling for endangered fish: conservation problem or conservation action? *Fish and Fisheries*. doi: 10.1111/faf.12076.
- Cowx IG. 1991. *Catch Effort Sampling Strategies: their Application in Freshwater Fisheries Management*. Fishing News Books, Blackwell: Oxford.
- De Graaf G, Bartley D, Jorgensen J, Marmulla G. in press. The scale of inland fisheries, can we do better? Alternative approaches for assessment. *Fisheries Management and Ecology*. doi: 10.1111/j.1365-2400.2011.00844.x.
- De Silva SS, Abery NW, Nguyen TTT. 2007. Endemic freshwater finfish of Asia: distribution and conservation status. *Diversity and Distributions* **13**: 172–184.
- Dhillon M. 2004. The mahseer of India Himalayas. *Rackelhanen Flyfishing Magazine*. Available at: www.rackelhanen.se/eng/10273.htm (accessed 16 April 2014).
- Dhu S. 1923. *The Angler in India or the Mighty Mahseer*. Natraj Publishers: Dehra Dun.
- Drenner RW, Gallo KL, Edwards CM, Rieger KE, Dibble ED. 1997. Common carp affect turbidity and angler catch rates of largemouth bass in ponds. *North American Journal of Fisheries Management* **17**: 1010–1013.
- Eschmeyer WN (ed). 2014. *Catalog of Fishes: genera, species, references*. <http://research.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>) Electronic version accessed 22 April 2014.
- Everard M. 2013. Safeguarding the provision of ecosystem services in catchment systems. *Integrated Environmental Assessment and Management* **9**: 252–259.
- Everard M, Kataria G. 2011. Recreational angling markets to advance the conservation of a reach of the Western Ramganga River, India. *Aquatic Conservation: Marine and Freshwater Ecosystems* **21**: 101–108.
- Gee AS, Milner NJ. 1980. Analysis of 70-year catch statistics for Atlantic salmon *Salmo salar* in the River Wye and implications for management of stocks. *Journal of Applied Ecology* **17**: 41–57.
- Granek EF, Madin EMP, Brown MA, Figueira W, Cameron DS, Hogan Z, Kristianson G, De Villiers P, Williams JE, Post J, *et al.* 2008. Engaging recreational fishers in management and conservation: global case studies. *Conservation Biology* **22**: 1125–1134.
- Inglis SD, Rosenau ML. 1994. Non-tidal sturgeon angler fishery of the lower Fraser River – angler card analysis. Ministry of Environment, Lands and Parks Regional Fisheries Report No. 241, Surrey, BC.
- IUCN. 2013. IUCN Red List of Threatened Species. Version 2013.2. [www.iucnredlist.org](http://www.iucnredlist.org). downloaded 22 April 2014
- Jensen OP, Gilroy DJ, Hogan Z, Allen BC, Hrabik TR, Weidel BC, Vander Zanden MJ. 2009. Evaluating recreational fisheries for an endangered species: a case study of taimen, *Hucho taimen*, in Mongolia. *Canadian Journal of Fisheries and Aquatic Sciences* **66**: 1707–1718.
- Knight JD, Rai A, D'Souza RKP. 2013. On the identities of *Barbus mussullah* Sykes and *Cyprinus curmuca* Hamilton with notes on the status of *Gobio canarensis* Jerdon (Teleostei: Cyprinidae). *Zootaxa* **3750**: 201–215.
- Kottelat M. 2013. The fishes of the inland waters of southeast Asia: a catalogue and core bibliography of the fishes known to occur in freshwaters, mangroves and estuaries. National University of Singapore.
- Langer RK, Ogale SN, Ayyappan S. 2001. *Mahseer in Indian Subcontinent - a Bibliography*. Central Institute of Fisheries Education (CIFE): Mumbai, India.
- Lehtonen H, Leskinen E, Sele' NR, Reinikainen M. 2009. Potential reasons for the changes in the abundance of pike, *Esox lucius*, in the western Gulf of Finland, 1939–2007. *Fisheries Management and Ecology* **16**: 484–491.
- MacDonald ASJ. 1948. *Circumventing the Mahseer*. Natraj Publishers: Dehra Dun.
- Mahon R. 1997. Does fisheries science serve the needs of managers of small stocks in developing countries. *Canadian Journal of Fisheries and Aquatic Sciences* **54**: 2207–2213.
- McMichael GA, Kaya CM. 1991. Relations among stream temperature, angling success for rainbow trout and brown trout, and fisherman satisfaction. *North American Journal of Fisheries Management* **11**: 190–199.
- Mezzerla M, Largiadèr CR. 2001. Evidence for selective angling of introduced trout and their hybrids in a stocked brown trout population. *Journal of Fish Biology* **59**: 287–301.
- Molur S, Smith KG, Daniel BA, Darwall WRT (compilers). 2011. The Status of Freshwater Biodiversity in the Western Ghats. International Union for Conservation of Nature (IUCN) Gland, Switzerland, & Zoo Outreach Organization (ZOO) Coimbatore, India.
- Nair S. 2010. Karnataka, *Where the Mahseer is Safe*. Available at: [www.deccanherald.com/content/77977/where-mahseer-safe.html](http://www.deccanherald.com/content/77977/where-mahseer-safe.html). (Accessed 16 April 2014).
- Nguyen TTT, Na-Nakorn U, Sukmanomon S, Ziming C. 2008. A study on phylogeny and biogeography of mahseer species (Pisces: Cyprinidae) using sequences of three mitochondrial DNA gene regions. *Molecular Phylogenetics and Evolution* **48**: 1223–1231.
- North E. 1980. The effects of water temperature and flow upon angling success in the River Severn. *Aquaculture Research* **11**: 1–9.
- North E, Hickley P. 1977. The effects of reservoir releases upon angling success in the River Severn. *Aquaculture Research* **8**: 86–91.
- Pinder AC. 2013. Conserving the mighty mahseer of South India. *FISH Magazine* **111**: 24–28.

- Pinder AC, Raghavan R. 2013. Conserving the endangered mahseers (*Tor* spp.) of India: the positive role of recreational fisheries. *Current Science* **104**: 1472–1475.
- Raghavan R. 2011. *Tor khudree*. In IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>. Downloaded on 22 April 2014.
- Raghavan R, Ali A. 2011. *Tor malabaricus*. In IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>. Downloaded on 22 April 2014.
- Raghavan R, Ali A, Dahanukar N, Rosser A. 2011. Is the Deccan Mahseer, *Tor khudree* (Sykes, 1839) (Pisces: Cyprinidae) fishery in the Western Ghats hotspot sustainable? A participatory approach to stock assessment. *Fisheries Research* **110**: 29–38.
- Reichard M, Jurajda P, Smith C. 2004. Spatial distribution of drifting cyprinid fishes in a shallow lowland river. *Archiv für Hydrobiologie* **159**: 395–407.
- Reid GM, MacBeath TC, Csatadi K. 2013. Global challenges in freshwater-fish conservation related to public aquariums and the aquarium industry. *International Zoo Yearbook* **47**: 6–45.
- Robinson AT, Clarkson RW, Forrest RE. 1998. Dispersal of larval fishes in a regulated river tributary. *Transactions of the American Fisheries Society* **127**: 772–786.
- Singh D, Sharma RC. 1998. Biodiversity, ecological status and conservation priority of the fish of the River Alaknanda, a parent stream of the River Ganges (India). *Aquatic Conservation: Marine and Freshwater Ecosystems* **8**: 761–772.
- Siraj SS, Christianus A, Kiat NC, de Silva SS (eds). 2007. Mahseer, the biology, culture and conservation. In *Proceedings of the International Symposium on the Mahseer*, 29–30 March 2006, Kuala Lumpur, Malaysia. Malaysian Fisheries Society: Serdang, Malaysia.
- Thomas HS. 1873. *The Rod in India*. 8vo: Mangalore.
- Trans World Fishing Team. 1984. *Quest for a Legendary Fish*. International Book Distributors: Dehra Dun.
- Ussi-Heikkila S, Wolter C, Klefoth T, Arlinghaus R. 2008. A behavioural perspective on fishing-induced evolution. *Trends in Ecology and Evolution* **23**: 419–421.